

BIROn - Birkbeck Institutional Research Online

Katus, Tobias and Eimer, Martin (2015) Lateralized delay period activity marks the focus of spatial attention in working memory: evidence from somatosensory event-related brain potentials. *Journal of Neuroscience* 35 (17), pp. 6689-6695. ISSN 0270-6474.

Downloaded from: <https://eprints.bbk.ac.uk/id/eprint/12038/>

Usage Guidelines:

Please refer to usage guidelines at <https://eprints.bbk.ac.uk/policies.html>
contact lib-eprints@bbk.ac.uk.

or alternatively

The Journal of Neuroscience

<http://jneurosci.msubmit.net>

JN-RM-5046-14R2

Lateralized delay period activity marks the focus of spatial attention in working memory: Evidence from somatosensory event-related brain potentials

Tobias Katus, Birkbeck College, University of London
Martin Eimer, Birkbeck College/University of London

Commercial Interest: No

Full Manuscript Title

**Lateralized delay period activity marks the focus of spatial
attention in working memory: Evidence from
somatosensory event-related brain potentials**

Running Title

Delay period activity marks the focus of attention

Tobias Katus^a & Martin Eimer^a

^aDepartment of Psychology, Birkbeck College, University of London, London
WC1E 7HX, United Kingdom.

Corresponding Author: Tobias Katus, Department of Psychology, Birkbeck College,
University of London, London WC1E 7HX, United Kingdom. Mail: t.katus@bbk.ac.uk

Number of Figures: 4

Total word count: 5392 (Abstract: 246, Introduction: 461; Materials & Methods: 1144;
Results: 734; Discussion: 1497; References: 788; Figure Legends: 520)

Conflict of Interest: The authors declare no competing financial interests.

20

21 *Acknowledgments:* This work was funded by the Deutsche Forschungsgemeinschaft
22 (DFG Grants KA 3843/1-1 and KA 3843/1-2), and supported by a grant from the
23 Economic and Social Research Council (ESRC), United Kingdom. We thank Anna
24 Grubert, Moran Aharoni and John Towler for constructive comments on the
25 manuscript, and Sue Nicholas for assistance in setting up the hardware for tactile
26 stimulation.

27

28 **Abstract**

29 The short-term retention of sensory information in working memory (WM) is known to
30 be associated with a sustained enhancement of neural activity. What remains
31 controversial is whether this neural trace indicates the sustained storage of
32 information, or the allocation of attention. To evaluate the storage and attention
33 accounts, we examined sustained tactile contralateral delay activity (tCDA
34 component) of the event-related potential (ERP). The tCDA manifests over
35 somatosensory cortex contralateral to task-relevant tactile information during stimulus
36 retention.

37 Two tactile sample sets (S1, S2) were presented sequentially, separated by
38 1.5 s. Each set comprised two stimuli, one per hand. Human participants memorized
39 the location of one task-relevant stimulus per sample set, and judged whether one of
40 these locations was stimulated again at memory test. The two relevant pulses were
41 unpredictably located on the same hand (stay trials) or on different hands (shift trials).
42 Initially, tCDA components emerged contralateral to the relevant S1 pulse. Sequential

loading of WM enhanced the tCDA after S2 was presented on stay trials. On shift trials, the tCDA's polarity reversed after S2 presentation, resulting in delay activity that was now contralateral to the task-relevant S2 pulse. The disappearance of a lateralized neural trace for the relevant S1 pulse did not impair memory accuracy for this stimulus on shift trials. These results contradict the storage account, and suggest that delay period activity indicates the sustained engagement of an attention-based rehearsal mechanism. In conclusion, somatosensory delay period activity marks the current focus of attention in tactile WM.

51

52 **Introduction**

Working memory (WM) allows for the sustained representation of information that is no longer perceptually present. Many WM tasks involve the retention of a specific stimulus attribute for comparison with a test stimulus, presented after a retention delay. Neural activity that persists during this delay is thought to reflect the sustained representation of information in memory (Wang, 2001; but see also Nairne, 2002; Sreenivasan et al., 2014). Sustained delay period activity has been found in prefrontal cortex (PFC; Fuster and Alexander, 1971; Romo and Salinas, 2003) and modality-specific sensory brain regions (touch: Kaas et al., 2013; Zhou and Fuster, 1996; vision: Sereno and Maunsell, 1998). Although elevated delay period activity is commonly observed in frontal and parietal areas, this activation may not directly reflect the retention of stimulus-specific information (e.g., Riggall and Postle, 2012), and could instead be linked to top-down attentional control aspects of WM tasks (Lewis-Peacock et al., 2012; LaRocque et al., 2013; Sreenivasan et al., 2014; Postle, 2015). The sustained representation of memorized features or objects is likely to be

67 implemented in sensory-perceptual brain areas (Curtis and D'Esposito, 2003;
68 D'Esposito, 2007; Emrich et al., 2013; Pasternak and Greenlee, 2005; Postle, 2006;
69 Jonides et al., 2005), even when these areas do not show sustained increases in
70 delay period activity that can be measured with fMRI (e.g. Harrison and Tong, 2009;
71 Riggall and Postle, 2012).

72 Event-related potential (ERP) studies of WM have revealed sustained delay
73 period activity with modality-specific neural generators. The tactile contralateral delay
74 activity (tCDA: Katus et al., 2014) and its visual counterpart (CDA: e.g. Vogel and
75 Machizawa, 2004) emerge when tactile or visual stimuli on one side are retained for
76 comparison with subsequent test stimuli as an enhanced negativity over
77 somatosensory or visual brain regions contralateral to the memorized stimulus set.
78 Although these components are usually interpreted as electrophysiological marker of
79 information storage in contralateral sensory areas (e.g., Vogel and Machizawa,
80 2004), they could also reflect a lateralized allocation of attention resources (van Dijk
81 et al., 2010).

82 In this study, we used the tCDA component to determine whether lateralized
83 somatosensory delay period activity reflects the retention of sensory information
84 (storage account) or the current focus of attention in WM (attention account). Two
85 bilateral tactile sample sets were presented sequentially. Each set involved a left-
86 and a right-hand pulse. Participants memorized the location of one pulse per set, and
87 judged whether one of these locations was stimulated again at memory test.
88 Critically, the two task-relevant pulses were unpredictably presented to the same
89 hand (stay trials) or to different hands (shift trials). If the tCDA component indicates
90 retention of tactile information in contralateral somatosensory cortex, it should
91 disappear on shift trials, where stimulus locations have to be simultaneously retained

on opposite hands. If it instead reflects the focus of attention in WM, the polarity of the tCDA should reverse on shift trials after the second sample set has been presented, due to the re-allocation of attention towards the most recently encoded item.

Methods

Participants

Brain activity was acquired from twelve neurologically unimpaired adult participants (mean age 32 years, range 25-41 years, 6 male, 9 right-handed). All participants gave informed written consent prior to testing. The study was conducted in accordance with the Declaration of Helsinki and approved by the Psychology Ethics Committee of Birkbeck College.

Stimuli and task design

Participants were seated in a dimly lit recording chamber with their hands covered from sight, viewing a monitor that showed a central white fixation cross against a black background. Eight mechanical tactile stimulators (four per hand) were attached to the distal phalanges of the index, middle, ring and small fingers of the left and right hands. Stimulators were driven by custom-built amplifiers using an eight-channel sound card (M-Audio, Delta 1010LT) controlled by MATLAB (MathWorks, Natick, MA). Continuous white noise masked sounds produced by tactile stimulation. All tactile stimuli were mechanical 100 Hz sinusoids (duration: 50 ms, intensity: 0.37 N).

The stimulation procedure involved two successive sets of bilaterally presented sample stimuli that were followed by a single test stimulus (see Figure 1A). The two sample sets (S1, S2) were separated by a 1.5 s delay, and the memory test stimulus followed S2 after additional 1.5 s. Each sample set consisted of a left-hand and a right-hand pulse. The pair of S1 pulses was simultaneously presented to one finger of the left and right hand, with left and right stimulus locations determined randomly and independently for each hand. The two S2 pulses were separated by an interstimulus interval (ISI) of 0.2 s. The order of S2 presentation (left-hand pulse preceding right-hand pulse, or vice versa; see Figure 1B) was randomly determined on each trial. The location of the two S2 pulses was randomly and independently selected, except that the two fingers that had already received an S1 pulse were not stimulated again. A unilateral memory test stimulus was presented 1.5 s after the first S2 stimulus to one finger of the left or right hand.

Participants had to memorize the locations of two cued sample pulses (one per sample set), and to decide whether one of the two memorized locations was stimulated again at memory test. Which tactile pulses were task-relevant was specified at the start of each block. Participants were instructed to remember the S1 pulse delivered to one of the two hands, and either the first or the second S2 pulse (which was equally likely to be presented to the same hand as the S1 pulse or to the other hand). The hand that was task-relevant for S1 (remember left-hand or right-hand S1 pulse) alternated between successive blocks. Six of the participants memorized left-hand S1 pulses in the first block, and the other six started the experiment by memorizing right-hand S1 pulses. The task-relevant temporal position of S2 (remember early or late S2 pulses) changed after six successive blocks, with six participants memorizing early S2 pulses in the first half of the experiment, and the

others memorizing late S2 pulses in their first six blocks. Unilateral test stimulus pulses were delivered with to one of the two fingers that had previously received a task-relevant S1 or S2 pulse (match trials, 50%) or to one of the other six fingers (mismatch trials, 50%). Participants were instructed to respond vocally ('a' for match trials, 'e' for mismatch trials) during the 1700 ms period after test stimulus onset, when a question mark replaced the fixation cross on the monitor. Vocal responses were recorded by a headset microphone. The next trial started after a random interval of 0.4-0.6 s after the end of this response period.

The experiment included 12 blocks with 40 trials each. One training block of 40 trials was run prior to the first experimental block. Another training block was run prior to the seventh experimental block, when task instructions regarding the temporal position of the task-relevant S2 pulse changed. Instructions stressed accuracy over speed and the need to avoid head and arm movements, and to maintain central gaze fixation. Feedback on task performance was provided on the computer screen after each experimental block.

insert Figure 1 about here

EEG data recording and analysis

EEG data were DC-recorded at 500 Hz from 64 active Ag/AgCl electrodes at standard locations of the extended 10-20 system, using a BrainVision DC amplifier. A

bipolar outer canthus montage (horizontal electrooculogram, HEOG) monitored lateral eye movements. Continuous EEG data were referenced to the left mastoid during recording, offline re-referenced to the arithmetic mean of both mastoids, and were submitted to a 40Hz low-pass finite impulse response filter (Blackman window, filter order 664). EEG epochs for the 3 s interval following the onset of the first sample set (S1) were corrected relative to a 200 ms pre-stimulus baseline.

Blind source separation of EEG data was performed with the Independent Component Analysis (ICA) algorithm provided by the EEGLab toolbox (Delorme and Makeig, 2004). Independent components related to stereotypical artifacts at anterior scalp regions (eye blinks, vertical and lateral eye movements) were identified by visual inspection (cf. Delorme et al., 2007) and subtracted from the EEG data. Lateral eye movements occurred on average on 5.6% of all trials, as indicated by a differential step function (step: 100 ms, threshold: 24 μ V), running on the bipolarized HEOG before ICA-based artifact correction. None of these epochs were marked by the same step function after EEG data had been corrected for lateral eye movements. Artifact rejection and the interpolation of noisy EEG channels was performed using *Fully Automated Statistical Thresholding for EEG Artifact Rejection* (FASTER; Nolan et al., 2010). 86.2% of all epochs were retained for statistical analyses (stay condition: 87.9%; shift condition: 84.5%), after artifact rejection and elimination of incorrect response trials.

ERPs from six electrodes at lateral central scalp regions (FC3/4, FC5/6, C3/4, C5/6, CP3/4, CP5/6) were separately averaged for ROIs contralateral and ipsilateral to the task-relevant S1 pulse. Statistical analyses were based on mean amplitudes of contra-/ipsilateral difference values for the S1-period (500-1500 ms after S1 onset) and the S2-period (500-1500 after S2 onset). In line with previous work (e.g. Katus et

al., 2014), the tCDA measurement time window for the S2-period started 300 ms after the potentially task-relevant late S2 pulse (which was presented 200 ms after the early S2 pulse). To ensure that measurement time windows were equally long for the S1- and S2-periods, the time window for the S1-period started 500 ms after the simultaneously presented S1 pulses. Data in spline-interpolated topographical voltage maps were collapsed across trials in which memory was required for the left- or right-hand pulse, by flipping electrode coordinates in left-hand memory trials over the midline. EEG data were collapsed across experimental blocks where the left- or right-hand S1 pulse was task-relevant, and blocks where the early or late S2 pulse was task-relevant, to focus on the critical comparison between stay and shift trials. Error bars in graphs showing difference values indicate 95% confidence intervals, which were calculated for each condition by t-tests against zero (i.e. no lateralized effect). Statistical significance of difference values is symbolized by asterisks (* for $p < 0.05$, ** for $p < 0.01$, *** for $p < 0.001$) and is marked by error bars (or colored shadings in the ERP plots) that do not overlap with the zero axis.

Results

Electrophysiological data

Figure 2 shows ERP waveforms for stay and shift trials during the 3 s interval following the onset of the first tactile sample set (S1). ERPs were averaged across lateral central electrodes (FC3/4, FC5/6, C3/4, C5/6, CP3/4, CP5/6) contralateral and ipsilateral to the task-relevant S1 pulse. The overall retention delay is divided into the S1-period (0.5-1.5 s after S1; memory load = 1 item) and the S2-period (0.5-1.5 s after S2; memory load = 2 items). Difference waveforms (Figure 2, bottom panel)

were calculated separately for stay and shift trials by subtracting ERPs ipsilateral to the task-relevant S1 stimulus from contralateral ERPs. Statistical analyses were conducted on mean amplitudes of these difference values in the S1- and S2-periods. Difference values that deviate significantly from zero indicate the presence of reliable lateralized effects.

insert Figure 2 about here

A sustained negativity (tCDA component) was present contralateral to the task-relevant S1 pulse in the S1-period, as indicated by difference values that were significantly different from zero in both stay and shift trials (stay trials: $t(11) = -5.174$, $p < 0.001$, average $-0.69 \mu V$; shift trials: $t(11) = -4.827$, $p = 0.001$, average $-0.67 \mu V$). Because the side of the task-relevant S2 pulse was unpredictable, tCDA amplitudes on stay and shift trials did not differ during the S1-period ($p > 0.7$). In the period after presentation of S2, tCDA amplitude further increased on stay trials, relative to the tCDA measured during the S1-period ($t(11) = -3.461$, $p = 0.005$). Critically, tCDA polarity reversed during the S2-period on shift trials, resulting in a statistically robust sustained negativity contralateral to the memorized S2 pulse in this period (test against zero: $t(11) = 3.472$, $p = 0.005$).

To avoid statistical comparisons of difference values with opposite signs (i.e. tCDA components with different polarities), analyses of the tCDA during the S2-

period were conducted on difference values that were calculated by subtracting ERPs ipsilateral to the task-relevant S2 stimulus from contralateral ERPs. Difference values were corrected relative to a 0.2 s baseline prior to S2 onset. The new baseline ensured that reliable lateralized effects triggered by the presentation of S2 (i.e., memory update effects) were marked by tCDA amplitude values that significantly differed from zero. As shown in Figure 3, robust tCDA components were found during the S2-period for stay trials ($t(11) = -7.082$, $p < 10^{-4}$) and shift trials ($t(11) = -7.954$, $p < 10^{-5}$). A repeated-measures ANOVA with the factors trial type (stay versus shift) and relevant S2 pulse (early versus late) revealed a highly significant main effect of trial type ($F(1, 11) = 20.013$, $p < 0.001$), and formally confirmed that the memory update effect on tCDA difference values was considerably larger in shift trials ($-1.24 \mu V$) relative to stay trials ($-0.56 \mu V$); see Figure 3. There were no tCDA differences between early and late pulses ($p > 0.6$).

To assess whether the tCDA components to S1 and S2 differed in size, we compared tCDA amplitudes in response to S1 (measured relative to the pre-S1 baseline) and to S2 (relative to a new pre-S2 baseline) on stay trials. The tCDA was numerically larger in the S1-period than in the S2-period ($-0.69 \mu V$ versus $-0.56 \mu V$), but this difference was not significant ($p > 0.3$).

insert Figure 3 about here

Behavioral performance

Participants responded correctly in 94.5% of all trials (stay trials: 96.8%, shift trials: 92.1%). Sensitivity indices (d') entered a three-way repeated measures ANOVA with the factors trial type (stay versus shift), relevant S1 pulse (left versus right hand), and relevant S2 pulse (early versus late); compare Figure 4A. A main effect of trial type showed that task performance was impaired on shift trials relative to stay trials ($F(1,11) = 19.439$, $p = 0.001$). No further effects or interactions were statistically reliable (all $ps > 0.3$).

The polarity of the tCDA component during the S2-period on shift trials was determined by the location of the memorized S2 pulse (see Figure 2). Seeing that, we examined whether the absence of delay period activity contralateral to the location of the task-relevant S1 pulse on these trials was linked to impaired memory accuracy for S1. Hit rates were calculated separately for trials where the test stimulus matched the location of the memorized S1 or S2 pulse (Figure 4B). A two-way repeated measures ANOVA with the factors tested item (S1, S2) and trial type (stay, shift) confirmed the reduced task performance for shift versus stay trials ($F(1,11) = 17.556$, $p = 0.002$), but did not reveal further statistically reliable effects or interactions (all $ps > 0.2$). Critically, hit rates on shift trials were not significantly reduced when memory was tested for S1 or S2 pulses (91.8% versus 92.8%; $p > 0.5$). Hence, the loss of delay period activity sensitive to the location of task-relevant S1 stimuli during the S2-period on shift trials was not accompanied by a selective impairment in retaining this information.

282 insert Figure 4 about here

283 -----

284 **Discussion**

285 The tactile contralateral delay activity (tCDA component) and its visual
286 counterpart (CDA component) both reflect different levels of neural activity between
287 hemispheres during the retention of tactile or visual information in WM. This
288 hemispherical asymmetry may directly reflect the storage of information in
289 contralateral sensory cortex (storage account; e.g. Harris et al., 2002), or
290 alternatively, the lateralized focus of spatial attention (attention account, e.g. van Dijk
291 et al., 2010). To dissociate these two accounts, we used a tactile memory matching
292 paradigm in which WM was sequentially loaded with two tactile stimuli, one per
293 sample set (S1, S2). Participants memorized the location of one pulse per sample
294 set, and decided whether any of these two locations was stimulated again at memory
295 test. The memorized stimuli were located on the same hand (stay condition), or on
296 different hands (shift condition), and tCDA components were measured during the
297 periods that followed the presentation of S1 and S2 pulses. For the S1-period, we
298 predicted a tCDA component over somatosensory cortex contralateral to the relevant
299 S1 pulse in both stay and shift trials. In the S2-period of shift trials, storage demands
300 were spatially balanced, because the relevant tactile stimuli had to be retained at
301 different hands. If the tCDA marks the sustained storage of task-relevant information
302 in contralateral somatosensory cortex, it should disappear during the S2-period of
303 shift trials. If delay period activity instead reflects the current focus of attention
304 (Lewis-Peacock et al., 2012; LaRocque et al., 2013; van Dijk et al., 2010), tCDA

components should emerge contralateral to the S2 pulse that was selected for memory update.

A sustained tCDA component was elicited over somatosensory cortex contralateral to the memorized S1 pulse during the S1-period (between 0.5 s and 1.5 s after S1 presentation), demonstrating that participants could successfully establish a lateralized memory representation of this tactile stimulus. This confirms observations from a previous tactile WM experiment where participants had to memorize either one or two tactile pulses delivered to one hand, while ignoring tactile stimuli presented simultaneously to the other hand (Katus et al., 2014). In this earlier study, reliable tCDA components were found for both WM load conditions, and tCDA amplitudes were larger when participants memorized two tactile stimuli rather than one stimulus on the same hand. Further evidence for the load sensitivity of the tCDA was obtained in the stay trials of the present experiment, even though tactile WM was now loaded sequentially, as the task-relevant S1 and S2 pulses were separated by a 1.5 s interval. The amplitude of the tCDA component on stay trials increased during the S2-period (between 0.5 s and 1.5 s after S2 onset) relative to the preceding S1-period (see Figure 2). Therefore, the sequential loading of WM with two tactile stimuli on the same hand enhances the contralateral delay activity similarly as when memory is required for two simultaneously presented stimuli (relative to memory for a single stimulus) (Katus et al., 2014).

The central new finding of the present study is that there was also a significant tCDA component during the S2-period on shift trials, contrary to the predictions of the storage account. Critically, this tCDA was triggered contralateral to the location of the task-relevant S2 pulse. On shift trials, a tCDA first emerged contralateral to the memorized S1 pulse during the S1-period. However, it changed polarity after the

task-relevant S2 pulse had been presented to the opposite hand (see Figure 2). In principle, this polarity reversal of the tCDA during the S2-period on shift trials could be explained if S2 would generally evoke larger tCDA components than S1. This possibility is ruled out by our observation that on stay trials, the tCDA elicited by S2 (after correction for a pre-S2 baseline) tended to be numerically smaller than the tCDA evoked by S1, although this difference was not statistically significant. The tCDA polarity reversal on shift trials therefore points towards a privileged state of information implicated in the most recent cognitive operation (cf. Zokaei et al., 2014; Postle et al., 2013). If the tCDA directly reflects memory storage, the presence of this component contralateral to the task-relevant S2 pulse would suggest that only this second stimulus was retained on shift trials, at the expense of the memory trace for the preceding S1 stimulus. However, this interpretation was not supported by behavioral data. If only the relevant S2 pulse was retained on shift trials, task performance should have been substantially impaired on trials where memory was tested for the relevant S1 pulse. Although performance was generally reduced for shift as compared to stay trials (Figure 4), there were no systematic performance differences when the location of the test stimulus matched with the relevant S1 or S2 pulse. Thus, both items were equally well retained on shift trials.

These findings strongly suggest that the representation of task-relevant information in tactile WM can be dissociated from a sustained modulation of neural activity in sensory regions, as indexed by the tCDA component. A similar conclusion has been drawn from recent studies of visual WM that employed multivariate pattern analysis (MVPA; Harrison and Tong, 2009; Serences et al., 2009) to decode the identity of memorized objects from fMRI (Lewis-Peacock et al., 2012) or EEG signals (LaRocque et al., 2013). In these studies, a retro-cue specified which of two visually presented sample stimuli would be relevant for an impending memory test. This test

was then followed by a second retro-cue and a second test. Even though the initially uncued stimulus had to be remembered because it could become relevant later, MVPA analyses did not detect an active neural trace for this unattended stimulus. A neural trace however emerged after this stimulus was marked as task-relevant by the second retro-cue. The observation that mnemonic content can be decoded from brain activity only while it is in the focus of attention suggests that fMRI and EEG measures are primarily sensitive to the attentional activation of stored information. Memory storage may be implemented by stimulus-specific changes in patterns of synaptic weights (e.g., Mongillo et al., 2008; Erickson et al., 2010), which would not lead to changes in brain activity that can be detected with fMRI or EEG methods (see Postle, 2015, for further discussion).

Our observation that the polarity of tCDA components changed between the S1- and S2-periods on shift trials, where task-relevant S1 and S2 pulses had to be retained on different hands, contradicts the storage account. It is however perfectly compatible with the hypothesis that the tDCA primarily reflects the momentary distribution of attention in somatotopic space (Katus et al., 2015). The net change of tCDA amplitudes between the S1- and S2-periods (memory update effect; see Figure 3) was twice as large on shift trials, where attention moved between hands, as compared to stay trials, where attention was re-allocated between two fingers on the same hand. This suggests that the sequential attentional selection of tactile locations on different body sides produces stronger changes in the relative activation of the two cerebral hemispheres than the sequential selection of two tactile locations on the same body side. The re-allocation of tactile attention between both hands may also account for the impaired performance on shift trials, as compared to stay trials. In a previous tactile dual-task study, a secondary perceptual attention task selectively impaired memory performance, when spatial attention had to be withdrawn from the

memorized location (Katus et al., 2012). Similar performance costs were found on shift trials in the present study. Finally, the task-relevant S1 and S2 locations were equally well retained on shift trials, although the relevant S1 pulse's location was not reflected by the tCDA component during the S2-period. This dissociation between behavioral and ERP data suggests that the sustained storage of information does not depend on an active neural trace (cf. Lewis-Peacock et al., 2012). Our results are furthermore consistent with a multi-component model of WM (Baddeley, 2003), which postulates distinct mechanisms for executive control and information storage.

The close link between the tCDA component and the allocation of spatial attention demonstrated here is in line with the idea that attention acts as a rehearsal mechanism in WM (Awh and Jonides, 2001; Awh et al., 2006), through the selective activation of mnemonic content that is currently relevant to behavioral goals (Lepsien and Nobre, 2006). Attended items in WM are thought to have a privileged state, relative to mnemonic content that is not relevant to ongoing cognitive operations (Cowan, 1997; Oberauer, 2009; Olivers et al., 2011). The attentional activation of stored information leads to modality-specific delay period activity (e.g. tCDA component), which marks the interaction between selection and storage mechanisms in sensory cortex. In this context, it is interesting to note that an fMRI study by Riggall and Postle (2012) found sustained delay period activity that was not stimulus-selective in frontal and parietal areas, whereas stimulus-specific information could be decoded from visual cortex using MVPA methods, in the absence of sustained activity enhancements in these posterior areas. These authors argued that sustained delay period activity reflects attentional control processes in higher-order cortex, and that stimulus-selective WM storage is based on distributed patterns of neural activation in sensory areas that can be detected with MVPA, but not with univariate

fMRI analyses. The present ERP results suggest that the maintenance of tactile representations is accompanied by a sustained modulation of neural activity in somatosensory cortex when focal attention is allocated to these representations. Unlike the sustained frontoparietal delay activity described by Riggall and Postle (2012), the tCDA component does not directly reflect attentional control processes themselves, but instead the effects of a flexible top-down attentional selection mechanism that modulates tactile WM representations in sensory cortex in a goal-directed fashion. The pattern of tCDA results observed in the present study therefore provides indirect evidence that sensory neurons contribute to the sustained storage of information in WM (sensory recruitment, cf. Jonides et al., 2005; Katus et al., 2014).

Conclusion

The dissociation between electrophysiological activity and memory accuracy in this study suggests that somatosensory delay period activity marks the attention-based rehearsal of information in tactile WM. The lateralization of tCDA components is not directly attributable to an asymmetric recruitment of the contra- versus ipsilateral hemispheres for the storage of somatosensory information in the brain, but reflects the spatially selective allocation of focal attention. Our findings also point towards a privileged state for information that was used to update an existing memory representation during the most recent attentional selection process.

References

430 Awh E, Jonides J (2001) Overlapping mechanisms of attention and spatial working
 431 memory. *Trends Cogn Sci* 5:119–126.

432 Awh E, Vogel EK, Oh SH (2006) Interactions between attention and working memory.
 433 *Neuroscience* 139:201–208.

434 Baddeley A (2003) Working memory: looking back and looking forward. *Nat Rev*
 435 *Neurosci* 4:829–839.

436 Cowan N (1997) Attention and memory. An integrated framework. New York, Oxford:
 437 Oxford University Press; Clarendon Press.

438 Curtis CE, D'Esposito M (2003) Persistent activity in the prefrontal cortex during
 439 working memory. *Trends Cogn Sci* 7:415–423.

440 Delorme A, Makeig S (2004) EEGLAB: an open source toolbox for analysis of single-
 441 trial EEG dynamics including independent component analysis. *J Neurosci*
 442 *Methods* 134:9–21.

443 Delorme A, Sejnowski T, Makeig S (2007) Enhanced detection of artifacts in EEG
 444 data using higher-order statistics and independent component analysis.
 445 *Neuroimage* 34:1443–1449.

446 D'Esposito M (2007) From cognitive to neural models of working memory. *Philos*
 447 *Trans R Soc Lond B Biol Sci* 362:761–772.

448 Emrich SM, Riggall AC, LaRocque JJ, Postle BR (2013) Distributed patterns of
 449 activity in sensory cortex reflect the precision of multiple items maintained in
 450 visual short-term memory. *J Neurosci* 33:6516–6523.

451 Erickson MA, Maramba LA, Lisman J (2010) A single brief burst induces GluR1-
 452 dependent associative short-term potentiation: a potential mechanism for short-
 453 term memory. *J Cogn Neurosci* 22:2530–2540.

454 Fuster JM, Alexander GE (1971) Neuron activity related to short-term memory.
 455 *Science* 173:652–654.

456 Harris JA, Miniussi C, Harris IM, Diamond ME (2002) Transient storage of a tactile
 457 memory trace in primary somatosensory cortex. *J Neurosci* 22:8720–8725.

458 Harrison SA, Tong F (2009) Decoding reveals the contents of visual working memory
 459 in early visual areas. *Nature* 458:632–635.

460 Jonides J, Lacey SC, Nee DE (2005) Processes of Working Memory in Mind and
 461 Brain. *Curr Dir Psychol Sci* 14:2–5.

462 Kaas AL, van Mier H, Visser M, Goebel R (2013) The neural substrate for working
 463 memory of tactile surface texture. *Hum Brain Mapp* 34:1148–1162.

464 Katus T, Andersen SK, Müller MM (2012) Common Mechanisms of Spatial Attention
 465 in Memory and Perception: A Tactile Dual-Task Study. *Cereb Cortex* 24:707–718.

466 Katus T, Grubert A, Eimer M (2014) Electrophysiological Evidence for a Sensory
 467 Recruitment Model of Somatosensory Working Memory. *Cereb Cortex*.

468 Katus T, Müller MM, Eimer M (2015) Sustained maintenance of somatotopic
 469 information in brain regions recruited by tactile working memory. *J Neurosci*
 470 35(4):1390-5.

471 LaRocque JJ, Lewis-Peacock JA, Drysdale AT, Oberauer K, Postle BR (2013)
 472 Decoding attended information in short-term memory: an EEG study. *J Cogn*
 473 *Neurosci* 25:127–142.

474 Lepsien J, Nobre AC (2006) Cognitive control of attention in the human brain.
 475 insights from orienting attention to mental representations. *Brain Res* 1105:20–31.

476 Lewis-Peacock JA, Drysdale AT, Oberauer K, Postle BR (2012) Neural evidence for
 477 a distinction between short-term memory and the focus of attention. *J Cogn*
 478 *Neurosci* 24:61–79.

479 Mongillo G, Barak O, Tsodyks M (2008) Synaptic theory of working memory. *Science*
 480 319:1543–1546.

481 Nairne JS (2002) Remembering over the short-term: the case against the standard
 482 model. *Annu Rev Psychol* 53:53–81.

483 Nolan H, Whelan R, Reilly RB (2010) FASTER: Fully Automated Statistical
 484 Thresholding for EEG artifact Rejection. *J Neurosci Methods* 192:152–162.

485 Oberauer K (2009) Design for a working memory. *Psychol Learn Motiv* 51:45–100.

486 Olivers, C. N. L., Peters J, Houtkamp R, Roelfsema PR (2011) Different states in
 487 visual working memory: when it guides attention and when it does not. *Trends*
 488 *Cogn Sci* 15:327–334.

489 Pasternak T, Greenlee MW (2005) Working memory in primate sensory systems. *Nat*
 490 *Rev Neurosci* 6:97–107.

491 Postle BR (2006) Working memory as an emergent property of the mind and brain.
 492 *Neuroscience* 139:23–38.

493 Postle BR (2015) The cognitive neuroscience of visual short-term memory. *Curr Opin*
 494 *Behav Sci* 1:40–46.

495 Postle BR, Awh E, Serences JT, Sutterer DW, D'Esposito M (2013) The positional-
 496 specificity effect reveals a passive-trace contribution to visual short-term memory.
 497 *PLoS ONE* 8:e83483.

498 Riggall AC, Postle BR (2012) The relationship between working memory storage and
 499 elevated activity as measured with functional magnetic resonance imaging. *J*
 500 *Neurosci* 32:12990–12998.

501 Romo R, Salinas E (2003) Flutter discrimination: neural codes, perception, memory
 502 and decision making. *Nat Rev Neurosci* 4:203–218.

503 Serences JT, Ester EF, Vogel EK, Awh E (2009) Stimulus-specific delay activity in
 504 human primary visual cortex. *Psychol Sci* 20:207–214.

505 Sereno AB, Maunsell JH (1998) Shape selectivity in primate lateral intraparietal
 506 cortex. *Nature* 395:500–503.

507 Sreenivasan KK, Curtis CE, D'Esposito M (2014) Revisiting the role of persistent
 508 neural activity during working memory. *Trends Cogn Sci* 18:82–89.

509 van Dijk H, van der Werf J, Mazaheri A, Medendorp WP, Jensen O (2010)
 510 Modulations in oscillatory activity with amplitude asymmetry can produce
 511 cognitively relevant event-related responses. *Proc Natl Acad Sci USA* 107:900–
 512 905.

513 Vogel EK, Machizawa MG (2004) Neural activity predicts individual differences in
 514 visual working memory capacity. *Nature* 428:748–751.

515 Wang XJ (2001) Synaptic reverberation underlying mnemonic persistent activity.
 516 *Trends Neurosci* 24:455–463.

517 Zhou YD, Fuster JM (1996) Mnemonic neuronal activity in somatosensory cortex.
 518 *Proc Natl Acad Sci USA* 93:10533–10537.

519 Zokaei N, Manohar S, Husain M, Feredoes E (2014) Causal evidence for a privileged
 520 working memory state in early visual cortex. *J Neurosci* 34:158–162.

521 **Figure Legends**

522 **Figure 1.** (A) Stimulation protocol. Two bilateral sample sets (S1, S2) were followed
 523 by one unilateral test stimulus. Each sample set involved two tactile pulses, one per
 524 hand, which were presented simultaneously for S1, and sequentially for S2. Only one
 525 pulse was task-relevant per sample set, and this was determined by spatial position
 526 for S1 (left or right hand) and temporal position for S2 (early or late pulse). (B)
 527 Experimental conditions, illustrated for blocks where participants had to remember
 528 the right-hand S1 pulse, and the early (top row) or late (bottom row) S2 pulse. The
 529 task-relevant sample stimuli (marked by black dots) were presented to the same
 530 hand on stay trials (left column), and to different hands on shift trials (right column).
 531 Stay and shift trials varied randomly and unpredictably within each block.



Participants' task was to judge whether one of the two memorized locations was stimulated again at memory test. Memory match trials (B1, B4) and mismatch trials (B2, B3) were equiprobable.

Figure 2. ERPs recorded over somatosensory scalp regions contralateral (bold line) and ipsilateral (thin line) to the memorized S1 pulse. Task-relevant S1 and S2 pulses were located on the same hand (green) on stay trials. On shift trials, they were located on different hands (red). Topographical difference maps show the scalp distribution of lateralized effects in the S1- and S2-periods in stay and shift trials. These maps represent the contralateral minus ipsilateral amplitude differences (defined relative to the side of the task-relevant S1 pulse). The bottom panel shows difference waves, obtained by subtracting ipsilateral ERPs from contralateral ERPs. Shaded areas represent 95% confidence intervals (CIs) for tests against zero (i.e. no lateralized effect). Time points when these shaded areas do not cross the x-axis ($y \neq 0$) indicate the presence of significant lateralized effects.


Figure 3. Memory update effects on tCDA amplitudes following the presentation of S2 pulses, relative to a 0.2 s baseline before S2 onset. The net change of tCDA amplitude during the S2-period was larger in shift relative to stay trials. The upper panel shows difference waveforms, calculated by subtracting ERPs ipsilateral to the task-relevant S2 pulse from contralateral ERPs. Shaded areas around the difference waveforms for stay (green) and shift trials (red) represent 95% CIs for tests of lateralized effects against zero. Difference maps illustrate the scalp distribution of lateralized effects in stay and shift trials. Bar graphs show mean tCDA amplitude

during the S2-period on stay and shift trials in blocks where the early or late S2 pulse was task-relevant. Error bars reflect 95% CIs for tests against zero.


Figure 4. (A) Sensitivity indices (d') for stay and shift trials, shown separately for blocks where the early or late S2 pulse was task-relevant. Performance was reduced on shift trials (white bars) relative to stay trials (black bars). (B) Hit rates on trials where the test stimulus matched the location of the task-relevant S1 or S2 pulse, shown separately for stay trials (black bars) and shift trials (white bars). Performance on shift trials was not impaired when the test stimulus matched the memorized S1 pulse relative to trials where it matched the S2 pulse.



Stay trials



Shift trials



Contralateral (bold line)





Ipsilateral (thin line)

to task-relevant S1 pulse


Stay
Shift

Difference waveform (contra minus ipsi)





A Sensitivity (d')



B Hit rates

